نموذج استرشادى (٣) لامتحان شهادة إتمام الدراسة الثانوية العامة ٢٠٢٥ / ٢٠٢٦ م

المادة : الإحصاء باللغة الإنجليزية (الشعبة الأدبية) الزمن : ثلاث ساعات

First: Multiple Choice questions (1 mark for each question)

(1) The scatter diagram represents a direct relationship between x and y is .....

(a) 

(b) 

(c) 

(d) 

(2) In the scatter diagram between the two variables x and y, if all the points belong to a straight line with a negative slope, then there is ..... correlation between x and y

(a)	nihilistic	(b)	weak inverse	(c)	perfect inverse	(d)	perfect direct

(3) The following data represents the scores of 11 students in a monthly test, represented in a stem-and-leaf plot, considering that the final score is 30. If the range of this data is equal to 19, then m = ...

Stem	Leaves
m	m m 4 5
2	m m 1 3
3	0 0 0

The key 3 | 0 represents 30

(a)	Zero	(b)	1	(c)	11	(d)	13

(4) The following box-plot illustrates the distribution of grades for a group of students in the mathematics examination:



then the semi interquartile range = ...

(a)	7	(b)	20	(c)	26	(d)	34

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(5)	If Z is a standard normal random variable, then P(|Z| < 1.53) = ...						
(a)	0.8740	(b)	0.4370	(c)	0.4115	(d)	0.4573

(6)	If A and B are mutually exclusive events from the sample space of a random experiment, and $P(A^{\setminus}) = 0.9$, then: P(A - B) = ...						
(a)	0.1	(b)	0.3	(c)	0.6	(d)	0.9

(7)	If A and B are two events from the sample space of a random experiment, and $P(A) = \frac{1}{3}$, $P(B) = \frac{3}{8}$, $P(A \cap B) = \frac{1}{4}$, then: $P(A^{\setminus} \cap B^{\setminus})$ = ...						
(a)	$\frac{11}{24}$	(b)	$\frac{13}{24}$	(c)	$\frac{3}{4}$	(d)	$\frac{1}{12}$

(8)	If μ is the mean of the normal random variable X, then $P(X \leq \mu)$ = ...						
(a)	0.1587	(b)	0.3413	(c)	0.5	(d)	0.8413

(9)	If X is a normal random variable with a mean μ = 50 and a standard deviation σ, given that P(X < 33) = 0.0446, then σ = ...						
(a)	10	(b)	20	(c)	30	(d)	50

(10)	The size of a sample is 144, and its standard deviation is 1.2, using a confidence of level 95%, then the estimation error equals.....						
(a)	0.12	(b)	2.96	(c)	1.96	(d)	0.196

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

Second: Multiple Choice questions (2 mark for each question)

(11)	To study the relationship between the variables $\mathcal{X}$ and y, if: $\sum D^2 = 8.5$, n = 6, then the spearman's correlation coefficient between the variables $\mathcal{X}$ and y $\simeq$						
(a)	0.76	(b)	1	(c)	0.24	(d)	zero

(12)	If the regression line equation is $\hat{y} = 8 + \frac{3}{4}x$, then the error in y when $x = 8$ if the tabulated value = 12 is						
(a)	2	(b)	3	(c)	5	(d)	7

(13)	If X is a random variable following a geometric distribution and P(X= 1) = 0.2, then the expectation = ...						
(a)	3	(b)	4	(c)	5	(d)	6

(14)	If the following data represents the weights of 10 gold items (measured in grams weight) if the data is represented by using the stems-and-leaves plot, then the third quartile = ...						
(a)	25	(b)	2.55	(c)	26.5	(d)	2.65

Stem	Leaves
1	4 5 6 7 8
2	3 4 5
3	1 1

The key 1|4 represents 1.4

(15)	If a coin is tossed two consecutive times and the upper face was observed, then the range of the random variable X, which represents the number of times heads appears, is						
(a)	{ 1 , 2 }	(b)	{ 0 , 1 , 2 }	(c)	{ 1 }	(d)	{ 2 }

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(16)	If A and B are events from the sample space of a random experiment, and if P(A) = 0.75, P(B) = 0.625, P(A ∪ B) = 0.875, then: $P(A^{\setminus} \mid B)$ = ...						
(a)	0.6	(b)	0.8	(c)	0.2	(d)	0.1

(17)	A study was carried out on a sample of female students regarding their pulse rate. Given that the sample size is 64, the standard deviation of the population of female students σ = 3.6, and the sample mean $\bar{x}$ = 18.4,then the confidence interval for the mean is ..... given that the confidence level is 95%.		
(a)	] 17.518 , 19.282 [	(b)	] 16.568 , 17.383 [
(c)	] 15.518 , 17.282 [	(d)	] 17.17 , 18.19 [

(18)	In studying the relationship between two variables x and y, if ∑ x = 316, ∑ y = 237, and n = 8, and the equation of the regression line of y on x is $\hat{y}$ = k x + 1.22, then the value of k ≃						
(a)	– 0.719	(b)	– 0.519	(c)	0.519	(d)	0.719

(19)	The following box plot illustrates the distribution of grades for a group of students in the statistics examination:  The semi-interquartile range = ...						
(a)	10	(b)	17.5	(c)	45	(d)	55

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(20)	If A and B are two independent events from the sample space of a random experiment, and P(A) = 0.4, P(B) = 0.25, then: P(A - B) = ...						
(a)	0.1	(b)	0.2	(c)	0.3	(d)	0.4

(21)	If A and B are two events from the sample space of a random experiment, and if P(A) =0.6, P(B) = 0.5, $P(A' \cup B') = 0.7$, then the probability of occurring at least one of the two events = ...						
(a)	0.3	(b)	0.4	(c)	0.5	(d)	0.8

(22)	A bag contains 7 blue balls and 3 red balls, if two balls are drawn one after the other without replacement, the probability that the first ball is blue and the second ball is red = ...						
(a)	$\frac{7}{33}$	(b)	$\frac{7}{30}$	(c)	$\frac{4}{33}$	(d)	$\frac{4}{11}$

(23) If the following box plot illustrates the area cultivated in acres in 25 villages in two different years:



Then the semi-interquartile range of the first year + the semi-interquartile range of the second year = ...

(a)	20	(b)	25	(c)	45	(d)	50

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(24)	If X is a random variable and the expectation = 3, $\sum x_r^2 \times f(x_r) = 14.5$, then the coefficient of variation ≃						
(a)	150.3%	(b)	150%	(c)	78%	(d)	183.3%

(25) If the number of hours 11 students spent using the internet weekly is as follows:

14, 35, 27, 21, 20, 40, 31, 18, 44, 40, 31

which of the following stem-and-leaf plots represents this data?

(a)

Stem	Leaves		
1	4	4	8
2	0	1	7
3	2	4	5
4	0	4	

The key 3 | 4 represents 34

(b)

Stem	Leaves		
1	4	8	
2	0	1	7
3	1	1	5
4	0	0	4

The key 3 | 5 represents 35

(c)

Stem	Leaves		
1	1	1	2
2	1	1	5
3	0	1	1
4	0	3	

The key 3 | 0 represents 30

(d)

Stem	Leaves		
1	1	1	2
2	0	1	1
3	2	4	8
4	0	3	

The key 3 | 8 represents 38

(26)	The size of a sample is 25, if its standard deviation is 10, using a confidence level of 95%, then the estimation error equals						
(a)	5.2	(b)	3.92	(c)	0.784	(d)	0.98

(27)	If the equation of the regression line is $\hat{y} = \frac{1}{2}x + 6$, then the expected value of y when $x = 4$ is						
(a)	2	(b)	4	(c)	6	(d)	8

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(28)	If the data represented in the stem-and-leaf plot represents the ages of 10 persons attending a club on a certain day, then: the lower quartile – the third quartile = ...						

Stem	Leaves
1	3 4
2	0 1 1 4 5 5
3	3 4

The key 2 \| 1 means 21

(a)	– 8.5	(b)	– 0.85	(c)	8.5	(d)	0.85

(29)	A box contains 5 red balls and 7 black balls. Two consecutive balls are drawn (without replacement),the probability that the second ball is red given that the first ball is black = ...						
(a)	$\frac{7}{12}$	(b)	$\frac{6}{11}$	(c)	$\frac{5}{11}$	(d)	$\frac{5}{12}$

(30)	If X is a random variable following a binomial distribution X ~ B (n, S) with an expectation of 8 and a variance of $\frac{20}{3}$, then the value of n = ...						
(a)	48	(b)	56	(c)	64	(d)	32

(31)	To study the relationship between the two variables x and y, if $\sum x = 160$, $\sum y = 170$, $\sum x^2 = 4882$, $\sum y^2 = 5244$, $\sum x\, y = 4993$, and n = 6, then the Pearson's correlation coefficient between the variables x and y $\simeq$						
(a)	– 0.98	(b)	– 0.89	(c)	0.891	(d)	0.896

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

(32)	If X is a continuous random variable and its probability density function is: $F(x) = \begin{cases} a\,x & , \quad 2 \leq x \leq 6 \\ zero & , \ otherwise \end{cases}$, then the value of a = ...

(a)	$\frac{1}{16}$	(b)	$\frac{1}{12}$	(c)	$\frac{1}{6}$	(d)	$\frac{1}{2}$

(33)	If X is a normally distributed random variable with mean μ and standard deviation σ, then P(X > μ + 0.8 σ) = ...

(a)	0.2881	(b)	0.2119	(c)	0.4641	(d)	0.7881

Third: Essay Questions "2 marks for each question"

(34)	If X is a discrete random variable with the probability distribution as follows : Calculate the value of m if the expectation is 3, then calculate the standard deviation of the random variable X.

x_r	1	2	4	m
$f(x_r)$	0.2	0.3	0.4	0.1

(35)	If the upper limit for the confidence interval for a sample mean is 15.98, the mean of the sample is 15, and the standard deviation of the sample is 16 at a confidence level of 95%, then find the sample size.

حقوق النشر محفوظة لوزارة التربية والتعليم والتعليم الفني 2025-2026

D.M.RAZK موقع الدكتور محمد رزق معلم الكيمياء التعليمي

موقع الدكتور محمد رزق التعليمي

امسح الباركود بالموبيل لتثبيت التطبيق وتحميل كل الملازم مجانا وبسرعه

MR

حمل التطبيق الأن

GET IT ON Google Play

ملازم ابتدائي | ملازم اعدادي | ملازم ثانـــوي | كورسات ونتائج امتحانات

موقع الدكتور محمد رزق
Dr.MoRizK

اجابه الامتحان الاسترشادي الثالث احصاء لغات انجليزي 2026 موقع الدكتور محمد رزق التعليمي

رقم السؤال	الاجـــــــابة
1	B
2	C
3	B
4	A
5	A
6	A

تم بواسطه موقع الدكتور محمد رزق التعليمي

7	B
8	C

موقع الدكتور محمد رزق معلم الكيمياء التعليمي D.M.RAZK

9	A
10	D
11	A
12	A
13	C
14	D
تم بواسطه موقع الدكتور محمد رزق التعليمي	
15	B
16	C
17	A
18	D
19	B
20	C
21	D

موقع الدكتور محمد رزق معلم الكيمياء التعليمي D.M.RAZK

22	B
23	C
24	C
25	B
تم بواسطه موقع الدكتور محمد رزق التعليمي	
26	B
27	D
28	A
29	C
30	A
31	D
32	A
33	B

موقع الدكتور محمد رزق معلم الكيمياء التعليمي D.M.RAZK



34





35

تم الاجابه من خلال موقع الدكتور محمد رزق التعليمي

والمصدر قناه مدرستنا 3

موقع الدكتور محمد رزق معلم الكيمياء التعليمي D.M.RAZK

موقع الدكتور محمد رزق التعليمي

امسح الباركود بالموبيل لتثبيت التطبيق وتحميل كل الملازم مجانا وبسرعه

MR

حمل التطبيق الأن

GET IT ON Google Play

ملازم ابتدائي | ملازم اعدادي | ملازم ثانـــوي | كورسات ونتائج امتحانات

موقع الدكتور محمد رزق
Dr.MoRizK